

<b>STN</b>	<b>Meranie rádioaktivity v prostredí. Vzduch-radón 222. Časť 5: Kontinuálna meracia metóda koncentrácie aktivity (ISO 11665-5: 2012).</b>	<b>STN EN ISO 11665-5</b>  40 4081
------------	---	--

Measurement of radioactivity in the environment -Air: radon-222 - Part 5: Continuous measurement method of the activity concentration (ISO 11665-5:2012)

Táto norma obsahuje anglickú verziu európskej normy.  
This standard includes the English version of the European Standard.

Táto norma bola oznámená vo Vestníku ÚNMS SR č. 01/16

Obsahuje: EN ISO 11665-5:2015, ISO 11665-5:2012

**122269**

---

Úrad pre normalizáciu, metrológiu a skúšobníctvo SR, 2016  
Podľa zákona č. 264/1999 Z. z. v znení neskorších predpisov sa môžu slovenské technické normy rozmnožovať a rozširovať iba so súhlasom Úradu pre normalizáciu, metrológiu a skúšobníctvo SR.

EUROPEAN STANDARD

**EN ISO 11665-5**

NORME EUROPÉENNE

EUROPÄISCHE NORM

September 2015

ICS 17.240

English Version

## Measurement of radioactivity in the environment -Air: radon-222 - Part 5: Continuous measurement method of the activity concentration (ISO 11665-5:2012)

Mesurage de la radioactivité dans l'environnement -  
Air: radon 222 - Partie 5: Méthode de mesure en  
continu de l'activité volumique (ISO 11665-5:2012)

Ermittlung der Radioaktivität in der Umwelt - Luft:  
Radon-222 - Teil 5: Kontinuierliches Messverfahren für  
die Aktivitätskonzentration (ISO 11665-5:2012)

This European Standard was approved by CEN on 12 June 2015.

CEN members are bound to comply with the CEN/CENELEC Internal Regulations which stipulate the conditions for giving this European Standard the status of a national standard without any alteration. Up-to-date lists and bibliographical references concerning such national standards may be obtained on application to the CEN-CENELEC Management Centre or to any CEN member.

This European Standard exists in three official versions (English, French, German). A version in any other language made by translation under the responsibility of a CEN member into its own language and notified to the CEN-CENELEC Management Centre has the same status as the official versions.

CEN members are the national standards bodies of Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, Former Yugoslav Republic of Macedonia, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and United Kingdom.



EUROPEAN COMMITTEE FOR STANDARDIZATION  
COMITÉ EUROPÉEN DE NORMALISATION  
EUROPÄISCHES KOMITEE FÜR NORMUNG

**CEN-CENELEC Management Centre: Avenue Marnix 17, B-1000 Brussels**

<b>Contents</b>	<b>Page</b>
European foreword.....	3

## European foreword

The text of ISO 11665-5:2012 has been prepared by Technical Committee ISO/TC 85 “Nuclear energy, nuclear technologies, and radiological protection” of the International Organization for Standardization (ISO) and has been taken over as EN ISO 11665-5:2015 by Technical Committee CEN/TC 430 “Nuclear energy, nuclear technologies, and radiological protection” the secretariat of which is held by AFNOR.

This European Standard shall be given the status of a national standard, either by publication of an identical text or by endorsement, at the latest by March 2016, and conflicting national standards shall be withdrawn at the latest by March 2016.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. CEN [and/or CENELEC] shall not be held responsible for identifying any or all such patent rights.

According to the CEN-CENELEC Internal Regulations, the national standards organizations of the following countries are bound to implement this European Standard: Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, Former Yugoslav Republic of Macedonia, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and the United Kingdom.

### Endorsement notice

The text of ISO 11665-5:2012 has been approved by CEN as EN ISO 11665-5:2015 without any modification.

---

---

**Measurement of radioactivity in the  
environment — Air: radon-222 —**

Part 6:  
**Spot measurement method of the activity  
concentration**

*Mesurage de la radioactivité dans l'environnement — Air: radon 222 —  
Partie 6: Méthode de mesure ponctuelle de l'activité volumique*





**COPYRIGHT PROTECTED DOCUMENT**

© ISO 2012

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office  
Case postale 56 • CH-1211 Geneva 20  
Tel. + 41 22 749 01 11  
Fax + 41 22 749 09 47  
E-mail [copyright@iso.org](mailto:copyright@iso.org)  
Web [www.iso.org](http://www.iso.org)

Published in Switzerland

<b>Contents</b>		Page
<b>Foreword</b> .....		iv
<b>Introduction</b> .....		v
<b>1 Scope</b> .....		1
<b>2 Normative references</b> .....		1
<b>3 Terms, definitions and symbols</b> .....		1
<b>3.1 Terms and definitions</b> .....		1
<b>3.2 Symbols</b> .....		2
<b>4 Principle</b> .....		2
<b>5 Equipment</b> .....		2
<b>6 Sampling</b> .....		3
<b>6.1 Sampling objective</b> .....		3
<b>6.2 Sampling characteristics</b> .....		3
<b>6.3 Sampling conditions</b> .....		3
<b>7 Detection</b> .....		3
<b>8 Measurement</b> .....		4
<b>8.1 Procedure</b> .....		4
<b>8.2 Influence quantities</b> .....		4
<b>8.3 Calibration</b> .....		4
<b>9 Expression of results</b> .....		4
<b>9.1 Radon activity concentration</b> .....		4
<b>9.2 Standard uncertainty</b> .....		5
<b>9.3 Decision threshold and detection limit</b> .....		5
<b>9.4 Limits of the confidence interval</b> .....		5
<b>10 Test report</b> .....		5
<b>Annex A (informative) Measurement method using scintillation cells</b> .....		7
<b>Bibliography</b> .....		13

## Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 2.

The main task of technical committees is to prepare International Standards. Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights.

ISO 11665-6 was prepared by Technical Committee ISO/TC 85, *Nuclear energy, nuclear technologies, and radiological protection*, Subcommittee SC 2, *Radiological protection*.

ISO 11665 consists of the following parts, under the general title *Measurement of radioactivity in the environment — Air: radon-222*:

- *Part 1: Origins of radon and its short-lived decay products and associated measurement methods*
- *Part 2: Integrated measurement method for determining average potential alpha energy concentration of its short-lived decay products*
- *Part 3: Spot measurement method of the potential alpha energy concentration of its short-lived decay products*
- *Part 4: Integrated measurement method for determining average activity concentration using passive sampling and delayed analysis*
- *Part 5: Continuous measurement method of the activity concentration*
- *Part 6: Spot measurement method of the activity concentration*
- *Part 7: Accumulation method for estimating surface exhalation rate*
- *Part 8: Methodologies for initial and additional investigations in buildings*

The following parts are under preparation:

- *Part 9: Method for determining exhalation rate of dense building materials*
- *Part 10: Determination of diffusion coefficient in waterproof materials using activity concentration measurement*



## Introduction

Radon isotopes 222, 220 and 219 are radioactive gases produced by the disintegration of radium isotopes 226, 224 and 223, which are decay products of uranium-238, thorium-232 and uranium-235 respectively, and are all found in the earth's crust. Solid elements, also radioactive, followed by stable lead are produced by radon disintegration<sup>[1]</sup>.

When disintegrating, radon emits alpha particles and generates solid decay products, which are also radioactive (polonium, bismuth, lead, etc.). The potential effects on human health of radon lie in its solid decay products rather than the gas itself. Whether or not they are attached to atmospheric aerosols, radon decay products can be inhaled and deposited in the bronchopulmonary tree to varying depths according to their size.

Radon is today considered to be the main source of human exposure to natural radiation. The UNSCEAR (2006) report<sup>[2]</sup> suggests that, at the worldwide level, radon accounts for around 52 % of global average exposure to natural radiation. The radiological impact of isotope 222 (48 %) is far more significant than isotope 220 (4 %), while isotope 219 is considered negligible. For this reason, references to radon in this part of ISO 11665 refer only to radon-222.

Radon activity concentration can vary by one to multiple orders of magnitude over time and space. Exposure to radon and its decay products varies tremendously from one area to another, as it depends firstly on the amount of radon emitted by the soil and the building materials in each area and, secondly, on the degree of containment and weather conditions in the areas where individuals are exposed.

The values commonly found in the continental environment are usually between a few becquerels per cubic metre and several thousand becquerels per cubic metre. Activity concentrations of one becquerel per cubic metre or less can be observed in the oceanic environment. Radon activity concentrations inside houses may vary from several tens of becquerels per cubic metre to several hundreds of becquerels per cubic metre<sup>[3]</sup>. Activity concentrations can reach several thousands of becquerels per cubic metre in very confined spaces.

The activity concentration of radon-222 in the atmosphere can be measured by spot, continuous and integrated measurement methods with active or passive air sampling (see ISO 11665-1). This part of ISO 11665 deals with radon-222 spot measurement methods.

**NOTE** The origin of radon-222 and its short-lived decay products in the atmospheric environment and other measurement methods are described generally in ISO 11665-1.

# Measurement of radioactivity in the environment — Air: radon-222 —

## Part 6: Spot measurement method of the activity concentration

### 1 Scope

This part of ISO 11665 describes radon-222 spot measurement methods. It gives indications for carrying out spot measurements, at the scale of a few minutes at a given place, of the radon activity concentration in open and confined atmospheres.

This measurement method is intended for rapid assessment of the radon activity concentration in the air. The result cannot be extrapolated to an annual estimate of the radon activity concentration. This type of measurement is therefore not applicable for assessment of the annual exposure.

The measurement method described is applicable to air samples with radon activity concentration greater than 50 Bq/m<sup>3</sup>.

**NOTE** For example, using an appropriate device, the radon activity concentration can be spot measured in the soil and at the interface of a material with the atmosphere (see also ISO 11665-7).

### 2 Normative references

The following referenced documents are indispensable for the application of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 11665-1, *Measurement of radioactivity in the environment — Air: radon-222 — Part 1: Origins of radon and its short-lived decay products and associated measurement methods*

ISO 11929, *Determination of the characteristic limits (decision threshold, detection limit and limits of the confidence interval) for measurements of ionizing radiation — Fundamentals and application*

ISO/IEC 17025, *General requirements for the competence of testing and calibration laboratories*

IEC 61577-1, *Radiation protection instrumentation — Radon and radon decay product measuring instruments — Part 1: General principles*

**koniec náhľadu – text ďalej pokračuje v platenej verzii STN**