

| | | |
|------------|---|--|
| STN | Meranie rádioaktivity v prostredí. Vzduch-radón 222. Časť 7: Akumulačná metóda stanovenia povrchovej exhalačnej rýchlosti (ISO 11665-7: 2012). | STN EN ISO 11665-7 40 4081 |
|------------|---|--|

Measurement of radioactivity in the environment - Air: radon-222 - Part 7: Accumulation method for estimating surface exhalation rate (ISO 11665-7:2012)

Táto norma obsahuje anglickú verziu európskej normy.
This standard includes the English version of the European Standard.

Táto norma bola oznámená vo Vestníku ÚNMS SR č. 01/16

Obsahuje: EN ISO 11665-7:2015, ISO 11665-7:2012

122271

Úrad pre normalizáciu, metrológiu a skúšobníctvo SR, 2016
Podľa zákona č. 264/1999 Z. z. v znení neskorších predpisov sa môžu slovenské technické normy rozmnožovať a rozširovať iba so súhlasom Úradu pre normalizáciu, metrológiu a skúšobníctvo SR.

EUROPEAN STANDARD

EN ISO 11665-7

NORME EUROPÉENNE

EUROPÄISCHE NORM

September 2015

ICS 17.240

English Version

Measurement of radioactivity in the environment - Air: radon-222 - Part 7: Accumulation method for estimating surface exhalation rate (ISO 11665-7:2012)

Mesurage de la radioactivité dans l'environnement -
Air: radon 222 - Partie 7: Méthode d'estimation du flux
surfaccique d'exhalation par la méthode d'accumulation
(ISO 11665-7:2012)

Ermittlung der Radioaktivität in der Umwelt - Luft:
Radon-222 - Teil 7: Anreicherungsverfahren zur
Abschätzung der Oberflächenexhalationsrate (ISO
11665-7:2012)

This European Standard was approved by CEN on 12 June 2015.

CEN members are bound to comply with the CEN/CENELEC Internal Regulations which stipulate the conditions for giving this European Standard the status of a national standard without any alteration. Up-to-date lists and bibliographical references concerning such national standards may be obtained on application to the CEN-CENELEC Management Centre or to any CEN member.

This European Standard exists in three official versions (English, French, German). A version in any other language made by translation under the responsibility of a CEN member into its own language and notified to the CEN-CENELEC Management Centre has the same status as the official versions.

CEN members are the national standards bodies of Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, Former Yugoslav Republic of Macedonia, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and United Kingdom.



EUROPEAN COMMITTEE FOR STANDARDIZATION
COMITÉ EUROPÉEN DE NORMALISATION
EUROPÄISCHES KOMITEE FÜR NORMUNG

CEN-CENELEC Management Centre: Avenue Marnix 17, B-1000 Brussels

| Contents | Page |
|------------------------|-------------|
| European foreword..... | 3 |

European foreword

The text of ISO 11665-7:2012 has been prepared by Technical Committee ISO/TC 85 “Nuclear energy, nuclear technologies, and radiological protection” of the International Organization for Standardization (ISO) and has been taken over as EN ISO 11665-7:2015 by Technical Committee CEN/TC 430 “Nuclear energy, nuclear technologies, and radiological protection” the secretariat of which is held by AFNOR.

This European Standard shall be given the status of a national standard, either by publication of an identical text or by endorsement, at the latest by March 2016, and conflicting national standards shall be withdrawn at the latest by March 2016.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. CEN [and/or CENELEC] shall not be held responsible for identifying any or all such patent rights.

According to the CEN-CENELEC Internal Regulations, the national standards organizations of the following countries are bound to implement this European Standard: Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, Former Yugoslav Republic of Macedonia, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and the United Kingdom.

Endorsement notice

The text of ISO 11665-7:2012 has been approved by CEN as EN ISO 11665-7:2015 without any modification.

**Measurement of radioactivity in the
environment — Air: radon-222 —**

**Part 7:
Accumulation method for estimating
surface exhalation rate**

Mesurage de la radioactivité dans l'environnement — Air: radon 222 —

Partie 7:

*Méthode d'estimation du flux surfacique d'exhalation par la méthode
d'accumulation*





COPYRIGHT PROTECTED DOCUMENT

© ISO 2012

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office
Case postale 56 • CH-1211 Geneva 20
Tel. + 41 22 749 01 11
Fax + 41 22 749 09 47
E-mail copyright@iso.org
Web www.iso.org

Published in Switzerland

| Contents | Page |
|--|-------------|
| Foreword | iv |
| Introduction | v |
| 1 Scope | 1 |
| 2 Normative references | 1 |
| 3 Terms, definitions and symbols | 1 |
| 3.1 Terms and definitions | 1 |
| 3.2 Symbols | 2 |
| 4 Principle of the measurement method for estimating surface exhalation rate | 2 |
| 5 Equipment | 4 |
| 6 Accumulation of radon in a container | 5 |
| 6.1 Accumulation characteristics | 5 |
| 6.2 Accumulation duration | 5 |
| 7 Sampling | 5 |
| 7.1 Sampling objective | 5 |
| 7.2 Sampling characteristics | 5 |
| 7.3 Sampling duration | 6 |
| 7.4 Volume of air sampled | 6 |
| 8 Detection method | 6 |
| 9 Measurement | 6 |
| 9.1 Procedure | 6 |
| 9.2 Influence quantities | 7 |
| 10 Expression of results | 7 |
| 10.1 Radon surface exhalation rate | 7 |
| 10.2 Standard uncertainty | 7 |
| 10.3 Decision threshold and detection limit | 7 |
| 10.4 Limits of the confidence interval | 8 |
| 11 Test report | 8 |
| Annex A (informative) Example of a sample results sheet | 10 |
| Annex B (informative) Estimation of radon surface exhalation rate using a continuous measurement method | 12 |
| Annex C (informative) Estimation of radon surface exhalation rate using a spot measurement method | 18 |
| Bibliography | 23 |

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 2.

The main task of technical committees is to prepare International Standards. Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights.

ISO 11665-7 was prepared by Technical Committee ISO/TC 85, *Nuclear energy, nuclear technologies, and radiological protection*, Subcommittee SC 2, *Radiological protection*.

ISO 11665 consists of the following parts, under the general title *Measurement of radioactivity in the environment — Air: radon-222*:

- *Part 1: Origins of radon and its short-lived decay products and associated measurement methods*
- *Part 2: Integrated measurement method for determining average potential alpha energy concentration of its short-lived decay products*
- *Part 3: Spot measurement method of the potential alpha energy concentration of its short-lived decay products*
- *Part 4: Integrated measurement method for determining average activity concentration using passive sampling and delayed analysis*
- *Part 5: Continuous measurement method of the activity concentration*
- *Part 6: Spot measurement method of the activity concentration*
- *Part 7: Accumulation method for estimating surface exhalation rate*
- *Part 8: Methodologies for initial and additional investigations in buildings*

The following parts are under preparation:

- *Part 9: Method for determining exhalation rate of dense building materials*
- *Part 10: Determination of diffusion coefficient in waterproof materials using activity concentration measurement*

Introduction

Radon isotopes 222, 220 and 219 are radioactive gases produced by the disintegration of radium isotopes 226, 224 and 223, which are decay products of uranium-238, thorium-232 and uranium-235 respectively, and are all found in the earth's crust. Solid elements, also radioactive, followed by stable lead are produced by radon disintegration^[1].

Radon is today considered to be the main source of human exposure to natural radiation. The UNSCEAR (2006) report^[2] suggests that, at the worldwide level, radon accounts for around 52 % of global average exposure to natural radiation. The radiological impact of isotope 222 (48 %) is far more significant than isotope 220 (4 %), while isotope 219 is considered negligible. For this reason, references to radon in this part of ISO 11665 refer only to radon-222.

The radon-222 half-life (3,8 days) is long enough for it to migrate from the rock producing it, through the soil, to the air^[3]. The radon atoms in the soil are produced by the disintegration of the radium-226 contained in the mineral grains in the medium. Some of these atoms reach the interstitial spaces between the grains: this is the phenomenon of emanation. Some of the atoms produced by emanation reach the soil's surface by diffusion and convection: this is the phenomenon of exhalation^{[3][4][5]}. These mechanisms are also brought into play in materials (building materials, walls, etc.).

The quantity of radon-222 reaching the open air per unit of time and per unit of surface is called the radon-222 surface exhalation rate and depends on the physical characteristics of the soil and weather conditions. When the ground is covered in snow or a layer of water, or is frozen, this surface exhalation rate can become very weak.

Values of the radon-222 surface exhalation rate observed in France, for example, vary between 1 mBq/m²/s and about 100 mBq/m²/s^{[6][7]}. In uranium-bearing ground, radon-222 surface exhalation rates in the order of 50 000 mBq/m²/s can be observed. By way of comparison, the United Nations Scientific Committee estimates the average surface exhalation rate on the surface of the globe at 20 mBq/m²/s^[8].

NOTE The origin of radon-222 and its short-lived decay products in the atmospheric environment and other measurement methods are described generally in ISO 11665-1.

Measurement of radioactivity in the environment — Air: radon-222 —

Part 7: Accumulation method for estimating surface exhalation rate

1 Scope

This part of ISO 11665 gives guidelines for estimating the radon-222 surface exhalation rate over a short period (a few hours), at a given place, at the interface of the medium (soil, rock, laid building material, walls, etc.) and the atmosphere. This estimation is based on measuring the radon activity concentration emanating from the surface under investigation and accumulated in a container of a known volume for a known duration.

This method is estimative only, as it is difficult to quantify the influence of many parameters in environmental conditions. This part of ISO 11665 is particularly applicable, however, in case of an investigation, a search for sources or a comparative study of exhalation rates at the same site. This part of ISO 11665 does not cover calibration conditions for the rate estimation devices.

The measurement method described is applicable for radon exhalation rates greater than 5 mBq/m²/s.

NOTE The uncertainty relating to the estimation of the result obtained by applying this part of ISO 11665 cannot guarantee that the true flux value is included in the uncertainty domain.

2 Normative references

The following referenced documents are indispensable for the application of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 11665-1, *Measurement of radioactivity in the environment — Air: radon-222 — Part 1: Origins of radon and its short-lived decay products and associated measurement methods*

ISO 11665-5, *Measurement of radioactivity in the environment — Air: radon-222 — Part 5: Continuous measurement method of the activity concentration*

ISO 11665-6, *Measurement of radioactivity in the environment — Air: radon-222 — Part 6: Spot measurement method of the activity concentration*

ISO/IEC 17025, *General requirements for the competence of testing and calibration laboratories*

IEC 61577-1, *Radiation protection instrumentation — Radon and radon decay product measuring instruments — Part 1: General principles*

koniec náhľadu – text ďalej pokračuje v platenej verzii STN