

STN	Poživatiny Stanovenie toxínov T-2 a HT-2 v obilninách a výrobkoch z obilnín pre dojčatá a malé deti metódou LC-MS/MS po prečistení metódou SPE	STN EN 16923 56 0526
------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------	----------------------------------------------

Foodstuffs - Determination of T-2 toxin and HT-2 toxin in cereals and cereal products for infants and young children by LC-MS/MS after SPE cleanup

Táto norma obsahuje anglickú verziu európskej normy.
This standard includes the English version of the European Standard.

Táto norma bola oznámená vo Vestníku ÚNMS SR č. 11/17

Obsahuje: EN 16923:2017

125520

Úrad pre normalizáciu, metrológiu a skúšobníctvo Slovenskej republiky, 2017
Podľa zákona č. 264/1999 Z. z. o technických požiadavkách na výrobky a o posudzovaní zhody a o zmene a doplnení niektorých zákonov v znení neskorších predpisov sa slovenská technická norma a časti slovenskej technickej normy môžu rozmnožovať alebo rozširovať len so súhlasom slovenského národného normalizačného orgánu.

EUROPEAN STANDARD

EN 16923

NORME EUROPÉENNE

EUROPÄISCHE NORM

May 2017

ICS 67.060; 67.230

English Version

Foodstuffs - Determination of T-2 toxin and HT-2 toxin in cereals and cereal products for infants and young children by LC-MS/MS after SPE cleanup

Produits alimentaires - Dosage des toxines T-2 et HT-2 dans les céréales et les produits céréaliers pour nourrissons et enfants en bas âge par CL-SM/SM après purification par SPE

Lebensmittel - Bestimmung von T-2-Toxin und HT-2-Toxin in Getreide und Säuglings- und Kleinkindernahrung auf Getreidebasis mit LC-MS/MS nach SPE-Reinigung

This European Standard was approved by CEN on 27 February 2017.

CEN members are bound to comply with the CEN/CENELEC Internal Regulations which stipulate the conditions for giving this European Standard the status of a national standard without any alteration. Up-to-date lists and bibliographical references concerning such national standards may be obtained on application to the CEN-CENELEC Management Centre or to any CEN member.

This European Standard exists in three official versions (English, French, German). A version in any other language made by translation under the responsibility of a CEN member into its own language and notified to the CEN-CENELEC Management Centre has the same status as the official versions.

CEN members are the national standards bodies of Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, Former Yugoslav Republic of Macedonia, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Serbia, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and United Kingdom.



EUROPEAN COMMITTEE FOR STANDARDIZATION
COMITÉ EUROPÉEN DE NORMALISATION
EUROPÄISCHES KOMITEE FÜR NORMUNG

CEN-CENELEC Management Centre: Avenue Marnix 17, B-1000 Brussels

Contents		Page
European foreword		3
Introduction		4
1	Scope	5
2	Normative references	5
3	Principle	5
4	Reagents	5
5	Apparatus and equipment	7
6	Procedure	9
7	Calculation	10
8	Precision	11
9	Test report	12
Annex A (informative) Example chromatograms (API 4000™)		13
Annex B (informative) Example conditions for suitable LC-MS/MS systems		17
Annex C (informative) Precision data		24
Bibliography		26

European foreword

This document (EN 16923:2017) has been prepared by Technical Committee CEN/TC 275 “Food analysis - Horizontal methods”, the secretariat of which is held by DIN.

This European Standard shall be given the status of a national standard, either by publication of an identical text or by endorsement, at the latest by November 2017, and conflicting national standards shall be withdrawn at the latest by November 2017.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. CEN shall not be held responsible for identifying any or all such patent rights.

This document has been prepared under a mandate given to CEN by the European Commission and the European Free Trade Association.

According to the CEN-CENELEC Internal Regulations, the national standards organizations of the following countries are bound to implement this European Standard: Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, Former Yugoslav Republic of Macedonia, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Serbia, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and the United Kingdom.

Introduction

The mycotoxin T-2 toxin and its metabolite HT-2 toxin belong to the group of trichothecenes which are produced by various *Fusarium* species. Cereals like maize, wheat, barley, oats, and rye are most likely to be affected.

WARNING 1 — Suitable precaution and protection measures need to be taken when carrying out working steps with harmful chemicals. The latest version of the hazardous substances ordinance, Regulation (EC) No 1907/2006 [3], should be taken into account as well as appropriate National statements e.g. such as in [4].

WARNING 2 — The use of this document can involve hazardous materials, operations and equipment. This document does not purport to address all the safety problems associated with its use. It is the responsibility of the user of this document to establish appropriate safety and health practices and determine the applicability of regulatory limitations prior to use.

WARNING 3 — T-2 toxin and its metabolite HT-2 toxin are known to have carcinogenic effects.

1 Scope

This European Standard describes a method for the determination of T-2 toxin and HT-2 toxin in cereals and cereal based products e.g. oats, intended for nutrition of infants and young children by high performance liquid chromatography (HPLC) coupled with tandem mass spectrometry (MS/MS) after cleanup by solid phase extraction (SPE) [5].

The method has been validated for HT-2 toxin in oat flour at levels of 9,3 µg/kg and 28,1 µg/kg, oat flakes at levels of 16,5 µg/kg and 21,4 µg/kg, and breakfast cereals (containing oat flakes) at a level of 8,1 µg/kg and for T-2 toxin in oat flour at levels of 4,4 µg/kg and 8,3 µg/kg, oat flakes at levels of 4,9 µg/kg and 6,6 µg/kg and breakfast cereals (containing oat flakes) at a level of 3,5 µg/kg.

Laboratory experiences [6] have shown that the method is also applicable to highly swelling materials (dry cereal based porridges and modified starches), but these were not examined in the method validation study. Details are outlined in 6.3.

The method can also be applied to oat-by-products at higher levels of T-2- and HT-2 toxin. In this case, the dilution steps need to be considered [6].

The method can also be applied to cereals and cereal products for infants and young children based on e.g. wheat, barley, and rice. In this case, the method needs to be in-house-validated for each material. At the time of the interlaboratory study, planned range was 10 µg/kg to 100 µg/kg, and it is known from the pre-study that the method works well in the whole range, although final validation was only done in the range from 3,5 µg/kg to 28,1 µg/kg.

2 Normative references

The following documents, in whole or in part, are normatively referenced in this document and are indispensable for its application. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

EN ISO 3696, *Water for analytical laboratory use - Specification and test methods (ISO 3696)*

koniec náhl'adu – text ďalej pokračuje v platenej verzii STN